



• **Name: Chang-Hwan Park**

• **Current Position & Affiliation:** Professor & Department of Internal  
Medicine, Chonnam National University  
Medical School

• **Country: South Korea**

• **Educational Background:**

1993: MD, Chonnam National University Medical School, Gwangju, Korea

2002: PhD, Chonnam National University Postgraduate School, Gwangju, Korea

• **Professional Experience:**

May 2003 – April 2004: Clinical Instructor

May 2004 - January 2005: Clinical Assistant Professor

February 2005 – March 2010: Assistant Professor

August 2008 – December 2009: Research Scholar, Geffen medical school, UCLA,  
California, USA

April 2010 – April 2015: Associate Professor

May 2015 – Present: Professor

• **Professional Organizations:**

The Korean Association of Internal Medicine

The Korean Society of Gastroenterology

Korean Society of Gastrointestinal Endoscopy

Korean Pancreatobiliary Association

• **Main Scientific Publications:**

(topic title, year, conference name or presenting books)

1. Current trends in the management of pancreatic cystic neoplasms in Korea: a national survey. Korean J Intern Med. 2022;37(1):63-72.
2. A National Survey on the Environment and Basic Techniques of Endoscopic Retrograde Cholangiopancreatography in Korea. Gut Liver. 2021;15(6):904-911.
3. Malignant potential of small pancreatic neuroendocrine neoplasm and its risk factors: A multicenter nationwide study. Pancreatology. 2021;21(1):208-214
4. Experience of Endoscopists in Endoscopic Retrograde Cholangiopancreatography in Surgically Altered Anatomy Patients. Clin Endosc. 2020;53(1):7-8.
5. A prospective, randomized, multicenter clinical trial comparing 25-gauge and 20-gauge biopsy needles for endoscopic ultrasound-guided sampling of solid pancreatic lesions. Surg Endosc. 2020;34(3):1310-1317.

6. Self-expandable metal stents for choledocholithiasis in Billroth II gastrectomy



- patients. *Hepatobiliary Pancreat Dis Int.* 2018;17(6):546-552.
7. Key lime (*Citrus aurantifolia*) inhibits the growth of triple drug resistant *Helicobacter pylori*. *Gut Pathog.* 2018;10:16.
  8. Restoration of common bile duct diameter within 2 weeks after endoscopic stone retraction is a preventive factor for stone recurrence. *Hepatobiliary Pancreat Dis Int.* 2018;17(3):251-256.
  9. Endoscopy. 2018 Apr;50(4):378-385. Aggressive intravenous hydration with lactated Ringer's solution for prevention of post-ERCP pancreatitis: a prospective randomized multicenter clinical trial. *Endoscopy.* 2018 Apr;50(4):378-385.
  10. Self-expandable metal stents for the extraction of common bile duct stones in patients receiving dual antiplatelet agents: a pilot study. *Surg Endosc.* 2018;32(2):1077-1081.